

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 16/28, C12N 5/20, C12Q 1/68,</b> <b>G01N 33/577, A61K 39/395, 49/00 //</b> <b>C07K 14/71</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/33772</b> <b>(43) International Publication Date:</b> 14 December 1995 (14.12.95)
<b>(21) International Application Number:</b> PCT/FI95/00337 <b>(22) International Filing Date:</b> 9 June 1995 (09.06.95) <b>(30) Priority Data:</b> 257,754 9 June 1994 (09.06.94) US <b>(71)(72) Applicants and Inventors:</b> ALITALO, Kari [FI/FI]; Nyyrikintie 4 A, FIN-02100 Espoo (FI). KAIPAINEN, Arja [FI/FI]; Messeniuksenkatu 7 A 9, FIN-00250 Helsinki (FI). KORHONEN, Jaana [FI/FI]; Agricolankatu 7 C 68, FIN-00530 Helsinki (FI). MUSTONEN, Tuija [FI/FI]; Pih- lajatie 8 A 2, FIN-00270 Helsinki (FI). PAJUSOLA, Katri [FI/FI]; Kasteholmantie 4 A 8, FIN-00900 Helsinki (FI). MATIKAINEN, Marja-Terttu [FI/FI]; Lankkistentanhua 12, FIN-23100 Mynämäki (FI). KARNANI, Päivi [FI/FI]; Mullintie 10 B 62, FIN-20300 Turku (FI). <b>(74) Agent:</b> ORION CORPORATION; Orion-Farmos, Research & Development, Patent Service, P.O. Box 65, FIN-02101 Espoo (FI).	<b>(81) Designated States:</b> AM, AT, AU, BG, BR, BY, CA, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KG, KP, KR, KZ, LT, LU, LV, MD, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, TJ, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
<b>(54) Title:</b> FLT4 RECEPTOR TYROSINE KINASE AND ITS USE IN DIAGNOSIS AND THERAPY		
<b>(57) Abstract</b>  Anti-FLT4 antibodies, especially monoclonal anti-FLT4 antibodies, which are useful as a specific marker for endothelial cells of lymphatic vessels and HEVs, as a diagnostic tool for detecting changes in lymphatic tissue, especially in lymphatic vessels and HEVs in disease states, such as lymphangioma, metastatic lymph nodes and inflammatory, infectious and immunological disease.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## FLT4 RECEPTOR TYROSINE KINASE AND ITS USE IN DIAGNOSIS AND THERAPY

### FIELD OF THE INVENTION

The present invention relates generally to receptor tyrosine kinases, nucleic acid probes and antibodies specifically recognizing such receptors, and the use of such probes and antibodies for identifying lymphatic vessels and high endothelial venules (HEV) in animal and human tissues and lymphatic endothelial cells in culture. More specifically the present invention is directed to FLT4, a receptor tyrosine kinase, to nucleic acid probes recognizing FLT4, to antibodies specific to FLT4 and to methods for identifying FLT4 expression in lymphatic vessels and ultimately diagnosing and treating disease states in animals and humans involving changes in lymphatic tissue, such as inflammatory, infectious and immunological diseases, metastatic lymph nodes and lymphangiomas.

### BACKGROUND OF THE INVENTION

The physiology of the vascular system, embryonic vasculogenesis and angiogenesis, blood clotting, wound healing and reproduction, as well as several diseases, involve the vascular endothelium lining the blood vessels. The development of the vascular tree occurs through angiogenesis and, according to some theories, the formation of the lymphatic system starts shortly after arterial and venous development by sprouting from veins (1,2).

After the fetal period endothelial cells proliferate very slowly, except during angiogenesis associated with neovascularization. Growth factors stimulating angiogenesis exert their effects via specific endothelial cell surface receptor tyrosine kinases.

The protein product of the FLT4 receptor tyrosine kinase cDNA, cloned from a human erythroleukemia cell line is N-glycosylated and contains seven immunoglobulin-like loops in its extracellular domain. The cytoplasmic tyrosine kinase domain of FLT4 is about 80 % identical at the amino acid

level with the corresponding domains of FLT1 and KDR and about 60 % identical with the receptors for platelet-derived growth factor, colony stimulating factor-1, stem cell factor and the FLT3 receptor <sup>(3)</sup>.

Although the biological function of FLT4 are as yet unknown, its  
5 restricted expression pattern indicated that its functions may involve the  
vascular endothelium. Our previous results revealed FLT4 mRNA expression  
in endothelial cells in developing vessels of several fetal organs as disclosed  
by Kaipainen et al., in *J. Exp. Med.* 178: 2077-2088, 1993. A comparison of  
FLT4, FLT1, and KDR/FLK-1 receptor mRNA signals showed overlapping,  
10 but distinct expression patterns in the tissues studied <sup>(4)</sup>. These data  
suggested that the receptor tyrosine kinases encoded by this gene family  
may have distinct functions in the regulation of the growth and/or  
differentiation of blood vessels.

A major function of the lymphatic system is to provide fluid return from  
15 tissues and transport many extravascular substances back to the blood. In  
addition, during the process of maturation, lymphocytes leave the blood,  
migrate through lymphoid organs and other tissues, and enter the lymphatic  
vessels, and return to the blood through the thoracic duct. Specialized  
venules, high endothelial venules, (HEVs) bind lymphocytes again and  
20 cause their extravasation into tissues. The lymphatic vessels and especially  
the lymph nodes thus play an important role in immunology and they are also  
sites of development of metastasis of different tumors.

Since the turn of the 20th century, three different theories concerning  
the embryonic origin of the lymphatic system have been presented.  
25 However, prior to the present invention, lymphatic vessels have been difficult  
to identify, because there are no specific markers available for them.

Lymphatic vessels are most commonly studied with the aid of lympho-  
graphy. In lymphography, X-ray contrast medium is injected directly into a  
lymphatic vessel. That contrast medium is distributed along the efferent  
30 drainage vessels of the lymphatic system. The contrast medium is collected  
in lymph nodes, where it stays for up to half a year, during which time X-ray  
analyses allow the follow-up of lymph node size and architecture. This

diagnostics is especially important in cancer patients with metastases in the lymph nodes and in lymphatic malignancies, such as lymphoma

### SUMMARY OF THE INVENTION

5 The present invention is directed to FLT4 peptides and other constructs and to the use of FLT4 as a specific marker for lymphatic endothelial cells.

The invention is also directed to nucleic acid probes and antibodies specifically recognizing FLT4, especially to monoclonal antibodies, and compositions containing such antibodies. Further disclosed in the present  
10 application is the use of such monoclonal antibodies for diagnostic purposes for detecting and measuring the amount of FLT4 receptors in tissues, especially in lymphatic tissues and in lymphatic endothelial cells.

In a preferred embodiment, the invention provides monoclonal antibodies specifically recognizing the FLT4 receptor. More specifically this  
15 invention provides a monoclonal antibody designated 9D9F9. The hybridoma cell line which produces monoclonal antibody 9D9F9 is deposited with the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) under the provisions of the Budapest Treaty (DSM accession number ACC2210).

20 Monoclonal antibodies labelled with a detectable marker are also provided. As used herein, the term detectable marker encompasses any detectable marker known to those skilled in the art. However, in a preferred embodiment of this invention, the detectable marker is selected from the group consisting of radioisotopes, fluorochromes, dyes, enzymes and biotin.  
25 For the purpose of this invention suitable radioisotopes include, but are not limited to <sup>125</sup>I and <sup>131</sup>I.

Monoclonal antibodies of the present invention may also be used in a method for detecting the presence of FLT4-receptors in a cell sample, comprising the steps of exposing a cell sample to a monoclonal antibody of  
30 the present invention and detecting the binding of said monoclonal antibody to FLT4-receptors.

Another aspect of the present invention thus relates to a method of determining the presence of FLT4-receptors in a cell sample, comprising the steps of:

- 5 (a) exposing a cell sample to a monoclonal antibody of the present invention;
- (b) detecting the binding of said monoclonal antibody to FLT4-receptors.

The exposure of a cell mixture to monoclonal antibodies of the invention can be in solution, as is the case for fluorescence-activated cell sorting, or it can be on solid tissue specimens, such as biopsy material, or it  
10 can be with the monoclonal antibody immobilized on a solid support, as is the case with column chromatography or direct immune adherence. The mixture of cells that is to be exposed to the monoclonal antibody can be any solution of blood cells or tissue cells. Preferably, the cell mixture is from  
15 normal or pathological tissue containing or suspected to contain lymphatic endothelial cells. After exposure of the cell mixture to the monoclonal antibody, those cells with FLT4 -receptors will bind to the monoclonal antibody to form an antibody-FLT4 -receptor complex. The presence of the antibody-FLT4 -receptor complex, and therefore FLT4 receptors, can be detected by methods known in the art. These methods include  
20 immunohistochemical methods standard in the art, such as immunofluorescence, FACS analysis, ELISA, IRMA (a sandwich type of immunochemistry assay), immunohisto-chemistry, RIA using <sup>125</sup>I-label and autoradiography.

The present invention also provides monoclonal antibodies  
25 conjugated to an imageable agent. As used herein, the term imageable agent includes, but is not limited to, radioisotopes. A preferred radioisotope is 99m-technetium.

In a specific embodiment, the invention is directed to a method for monitoring lymphatic vessels and their endothelial cells in tissue samples  
30 and in organisms. The present invention further provides clinical detection methods describing the state of lymphatic tissue, and especially lymphatic vessels (inflammation, infection, traumas, growth, neoplasia etc.) and methods for detecting lymphatic vessels and thus lymphatic vascularization in an organism.

More specifically the present invention provides a method for detecting and identifying lymphatic changes characterized by FLT4 expression in connection to metastatic cancers, inflammatory, infectious and immunological conditions, which method comprises the steps of

5 (a) obtaining a tissue and/or body fluid sample suspected of lymphatic changes, and

(b) contacting said sample with a FLT4-specific monoclonal antibody under conditions suitable for forming a complex between the monoclonal antibody and the antigen, and

10 (c) detecting the presence of any complex formed.

A tissue which may be detected by this method is any normal, precancerous or cancerous solid tumor tissue with FLT4-containing lymphatic cells or cells which express the FLT4-receptor. In one embodiment of the present invention, the monoclonal antibody is labelled with a  
15 detectable marker as described herein. Methods of the invention are useful for detecting and differentiating various forms of cancer, especially metastases in the lymph nodes and other lymphatic malignancies, such as lymphomas, as well as lymphangiosarcomas.

A method of imaging the presence of lymphatic vessels, high  
20 endothelial venules or lymph nodes in human patients, is also provided by this invention. This method comprises administration of labelled antibodies and detection by imaging at sites where FLT4 expressing cells are present, in lymphatic vessels or lymph nodes.

The invention is further directed to a method of stimulating or  
25 antagonizing the function of FLT4 in lymphatic vascularization and in inflammatory, infectious and immunological conditions, said method comprising inhibiting the FLT4-mediated lymphatic vascularization by providing amounts of a FLT4-binding compound sufficient to block the FLT4  
endothelial cell sites participating in such reaction, especially where FLT4  
30 function is associated with a disease such as metastatic cancers, lymphomas, inflammation (chronic or acute), infections and immunological diseases.

**BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1. Expression of FLT4 mRNA in mouse tissues A. Hybridization of polyadenylated RNA isolated from the indicated tissues of adult mice. The size of the FLT4 mRNA band is given in kilobases. B. RNase protection analysis of RNA isolated from mouse embryos of various gestational ages (E8-E18) and from a newborn mouse (1 day). Sample E8+P contains also the placenta. The size of the probe and the protected FLT4 fragment are given in nt;  $\beta$ -actin was used as a control.

Figure 2. Expression of FLT4 mRNA in a 7.5-, 8.5- and 11.5-day p.c. embryos. Darkfield and brightfield photomicrographs of *in situ* autoradiograms are shown. No expression of FLT4 mRNA could be detected in a 7.5-day embryo (A). The FLT4 expression of an 8.5-day p.c. mouse embryo is shown in (B) and (C). Arrows point to FLT4 positive cells in the endothelium of posterior cardinal vein (cv), in the allantois (al) in (B) and in angioblasts (ab) of the head mesenchyme in (C). In a 8,5-day p.c. placenta FLT4 transcripts can be seen in endothelial cells of venous lacunae (vl) (D). Panels E and F show a comparison of FLT4 and Tie hybridization signals in 11.5-day p.c. embryos. The region of the developing dorsal aorta and metanephros (mn) is shown (20x). Note that the dorsal aorta is negative for FLT4, but positive for Tie mRNA, whereas both probes hybridize with the endothelium of the subendocardial vein (sv). Also, the FLT4 probe gives a signal from the metanephric vein (v), whereas Tie mostly hybridizes with the metanephric capillaries (c, arrows). da: dorsal aorta, ng: neural groove. Scale bar: 30  $\mu$ m.

Figure 3. FLT4 mRNA expression in a 12.5-day p.c. embryo. A sagittal section through the axillar plane is shown. Note that FLT4 mRNA is prominent in dilated vessels of the axilla (ax), in a plexus-like pattern in the periorbital (po) region, in the paravertebral tissue (arrowheads) and in the subcutaneous (sc) tissue. b: brain, li: liver. Scale bar: 5  $\mu$ m.

Figure 4. FLT4 in 14 and 16.5 day p.c. embryos. Panels A and B show bright and darkfield images of a midsagittal section. po: periorbital region, lj: lower jaw, ne: neck region, sc: subcutis, mt, mesenterium, ao, aorta. dt: thoracic duct. (C) shows a transverse section of a 16.5 day embryo in



hematoxylin-eosin staining. th: thymus, tr: trachea, e: esophagus, ca: carotid artery, ba: brachiocephalic artery, dt: thoracic duct. (D) shows a higher magnification (40x) of the region of ductus thoracicus; the autoradiographic grains can be seen over the endothelial cells. Also, the small vessel (v) in the upper part of the photograph is positive. Scale bar: 10  $\mu$ m (A-C), 1  $\mu$ m (D).

Figure 5. Comparison of FLT4 and Tie mRNA expression in cultured endothelial cells. Northern blot analysis of polyadenylated RNA from human foreskin microvascular (MV), femoral vein (VE), aortic (AO) and umbilical vein (HU) endothelial cells. For a comparison, the hybridization signal of the Tie receptor tyrosine kinase mRNA is shown. The bands resulting from the unspecific binding of the probe to the ribosomal RNA are marked with asterixes.

Figure 6. FLT4 in adult human lymphatic vessels of the mesenterium (A,B), lung (C,D), and tonsil (E,F). Note that only the lymphatic vessels in A and C give a FLT4 signal, whereas the veins, capillaries and the arteries are negative for FLT4 mRNA. In the tonsil, the signal is found in the endothelia of some HEVs. Scale bar: 200  $\mu$ m.

Figure 7. FLT4 mRNA in normal (A, B) and metastatic (C, D) lymph node and in lymphangioma (E-G). Arrowheads mark the lymphatic sinuses and HEVs, which are FLT4 positive. A comparison of FLT4 and von Willebrand factor signals shows both in the lymphatic endothelium, but only von Willebrand factor signal in the capillary (c) and venous (v) endothelia. Scale bars: 10  $\mu$ m (A-D) and 100  $\mu$ m (E-G).

Figure 8. FLT4 expression in fetal mesenterial vessels detected by immunoperoxidase staining. Sections were stained with affinity-purified anti-FLT4 antibodies (A), with antigen-blocked antiserum (B) and with preimmune serum (C) and with antiserum specific against the factor VIII-related antigen (D). Note that staining is confined to some, but not all vessels (v). Scale bar, 0.05 mm.

### DETAILED DESCRIPTION OF THE INVENTION

Recognizing the importance of identifying changes in lymphatic tissues, especially in lymph nodes in connection to metastatic cancers and immunological disease states, the present inventors have shown that FLT4 is  
5 a specific marker that detects lymphatic vessel endothelium and therefore useful as a marker for lymphatic changes in pathological states in man.

The present inventors have earlier shown that the expression pattern of FLT4 in comparison to FLT1 and KDR differs greatly in tissues of 18-week-old human fetuses <sup>(4)</sup>. In order to understand the role of FLT4 during  
10 development, the inventors cloned partial cDNAs for mouse FLT4. Using these probes in *in situ* hybridization, FLT4 mRNA expression during mouse development was analysed and it was found that FLT4 is expressed during vasculogenesis and angiogenesis of the lymphatic system. The relevance of these findings was also confirmed in normal and pathological human adult  
15 tissues, as FLT4 was found in lymphatic endothelial cells of human adult tissues both in normal and pathological conditions, as well as in some high endothelial venules (HEVs).

The cloning of mouse FLT4 cDNA fragments showed that their deduced amino acid sequence is almost identical with the corresponding  
20 human sequence (amino acid identity about 96 % in both segments studied). Further evidence for the identity of the mouse FLT4 cDNA was obtained from Northern hybridization where probes from both species yielded the typical 5.8 kb mRNA signal from mouse tissues. Analysis of RNA isolated from various tissues of adult mice showed FLT4 expression in the liver, lung,  
25 heart, spleen and kidney, with no or very little hybridization in the brain and testes. This pattern is similar to the pattern reported earlier by Galland et al. <sup>(5)</sup>. The results of RNase protection suggested that the FLT4 gene is needed during mouse development, starting from 8.5 day p.c. embryos, and the relative expression levels appeared quite stable.

30 For the *in situ* hybridization two fragments of mouse FLT4 cDNA were selected, which encode sequences of the extracellular domain. This allowed a clear distinction of the hybridization pattern from the related FLK-1 and FLT-

1 receptor patterns, which show only a very low degree of sequence identity with FLT4 in the extracellular region (6, 7, 8, 9). FLT4, similarly to the FLK-1, FLT-1, Tie and Tek endothelial receptor tyrosine kinase genes was not expressed in 7.5 day p.c. embryos. In a 8.5-day p.c. embryo the strongest  
5 FLT4 signals were localised in the allantois, the angioblasts of head mesenchyme and the cardinal vein. In contrast, the the dorsal aorta, endocardium of the heart and angioblasts of the yolk sac were negative, unlike for Tie, Tek, FLK-1 and FLT-1, Tie and Tek (10, 8). The restriction of FLT4 expression to the venous system was even more clear in samples from  
10 11.5 day mouse embryos; where the Tie mRNA was expressed also in arteries. In 12.5-day p.c. embryos the FLT4 signal decorated developing venous and presumptive lymphatic endothelia, but unlike for the endothelial Tie receptor tyrosine kinase, arterial endothelia were negative. During later stages of development FLT4 mRNA became restricted to vascular plexuses  
15 devoid of blood cells, representing developing lymphatic vessels. Only the lymphatic endothelium and some high endothelial venules expressed FLT4 mRNA in adult human tissues. Increased expression occurred in lymphatic sinuses and high endothelial venules in metastatic lymph nodes and in lymphangioma.

20 Due to difficulties in the interpretation of data from mouse embryos, human endothelia were studied, because the lymphatic system is much better defined in humans. Also, cells established from various endothelia could be studied in cell culture to see if the specificity of FLT4 expression persists in *in vitro* conditions. Endothelial cells lines are known to lose  
25 differentiated features upon in vitro culture. Therefore, it was not unexpected that they were negative for FLT4. Cultured aortic endothelial cells were also devoid of FLT4 mRNA. However, signals were obtained from human endothelial cells grown from the microvasculature and from femoral and umbilical veins. Thus, at least some of the specificity of FLT4 expression was  
30 retained in cell culture.

*In situ* hybridization analysis of adult human tissues confirmed the restriction of FLT4 to the lymphatic system seen in the developing mouse embryos. FLT4 expression was seen in the lymphatic endothelia and in the sinuses of human lymph nodes. Interestingly, also some of the HEVs, which  
35 have a cuboidal endothelium, shown to function in the trafficking of

leukocytes to the lymph nodes, were FLT4 positive. Furthermore, a parallel hybridization analysis showed that FLT4 mRNA levels were enhanced in these structures in metastatic as compared to normal lymph nodes. FLT4 was also very prominent in lymphangiomas, which are benign tumours composed  
5 of connective tissue stroma and growing, endothelial-lined lymphatic channels. FLT4 mRNA was restricted to the lymphatic endothelium of these tumors and absent from their arteries, veins and capillaries. In the human lung we were able to identify lymphatic structures, which were the only FLT4 positive vessels in this tissue.

10 The foregoing results suggest that FLT4 is a novel marker for lymphatic vessels and some high endothelial venules in human adult tissues. They also support the theory on the venous origin of lymphatic vessels. FLT4, as a growth factor receptor, may be involved in the differentiation and functions of these vessels.

15 These results, combined with the FLT4-binding compounds according to the present invention, allows a selective labelling of lymphatic endothelium, especially by using antibodies of the present invention coupled to radioactive, electron-dense or other reporter substances, which can be visualized. It may be possible to inject into the lymphatic system substances,  
20 containing FLT4 receptor internalization-inducing monoclonal antibodies, and thereby transport predefined molecules into the lymphatic endothelium. Also, it may be possible to use such the FLT4-binding compounds according to the invention for the detection of high endothelial venules, especially activated HEVs, which express enhanced levels of the FLT4 receptor. To our  
25 knowledge, no such specific markers are currently available for lymphatic endothelium.

The following examples are given merely to illustrate the present invention and not in any way to limit its scope.

## EXAMPLES

### EXAMPLE 1

#### Cloning of mouse FLT4 cDNA probes

Approximately  $10^6$  plaques from a iFIX®II genomic library from 129SV mice (Stratagene) was screened with the S2.5 human FLT4 receptor cDNA fragment covering the extracellular domain <sup>(3)</sup>. A 2.5 kb Bam HI fragment was subcloned from a positive plaque and sequenced from both ends. From this subclone, polymerase chain reaction was used to amplify and clone into the pBluescript KSII+/- vector (Stratagene) an exon fragment covering nucleotides 1745-2049 of the mouse FLT4 cDNA sequence <sup>(9)</sup>.

A second fragment covering nucleotides 1-192 was similarly cloned.

### EXAMPLE 2

#### Analysis of FLT4 mRNA in mouse tissues

Total RNA was isolated from developing embryos (8-18 days p.c. and one day old mice) according to Chomczynski et al. <sup>(11)</sup>. The sample from 8 day p.c. embryos included also the placenta.

For RNase protection analysis, RNA probe was generated from the linearized FLT4 plasmid obtained according to Example 1 using [<sup>32</sup>P]-UTP and T7 polymerase for the antisense orientation. The  $\beta$ -actin probe used corresponds to nucleotides 1188-1279 of the published mouse  $\beta$ -actin sequence <sup>(12)</sup>. After purification in a 6% polyacrylamide/7M urea gel, the labelled transcripts were hybridized to 30  $\mu$ g of total RNA overnight at 52 °C. Unhybridized RNA was digested with RNase A (10 U/ml) and T1 (1  $\mu$ g/ml) at 37 °C, pH 7.5 for 1 h. The RNases were inactivated by proteinase K digestion at 37 °C for 15 min and the samples were analysed in a 6% polyacrylamide/7M urea gel.

The pattern of expression of FLT4 analysed in this experiment showed that very weak mRNA signals were obtained from lung, liver, heart, kidney, skeletal muscle and spleen, whereas testis and brain were apparently

without specific signal (Fig. 1A). Analysis of a series of RNAs collected during different phases of mouse development by RNase protection assay showed that the FLT4 mRNA was expressed throughout embryogenesis from day 8 p.c. to newborn mice without great variations in signal intensity (Fig. 1B).

### EXAMPLE 3

#### *In situ* hybridization for FLT4 in mouse embryos

To better assign FLT4 transcripts to cells and tissues, sections of 7.5 and 8.5 day p.c. mouse embryos were hybridized with labelled FLT4 RNAs. Mouse embryos were derived from matings of CBA and NMRI mice. Pregnant mice were killed by cervical dislocation and the embryos were either immediately frozen or transferred via phosphate buffered saline into 4% paraformaldehyde. The embryos and isolated mouse organs were fixed for 18 h at 4°C, dehydrated, embedded in paraffin and cut into 6 µm sections.

RNA probes (antisense and sense) of 192 and 349 nucleotides (see Example 1) were generated from linearized plasmids using [<sup>35</sup>S]-UTP. *In situ* hybridization of sections was performed according to Wilkinson et al. (13, 14), incorporated by reference herein, with the following modifications: 1) instead of toluene, xylene was used before embedding in paraffin wax, 2) 6 µm sections were cut, placed on a layer of diethyl pyrocarbonate-treated water on the surface of glass slides pretreated with 2% 3-triethoxysilylpropylamine, 3) alkaline hydrolysis of the probes was omitted, and 4) the high stringency wash was for 80 min at 65°C in a solution containing 30 mM DTT and 1 x SSC. The sections were covered with NTB-2 emulsion (Kodak) and stored at 4°C. The slides were exposed for 14 days, developed and stained with hematoxylin. Control hybridizations with sense strand and RNase A-treated sections did not give a specific signal above background.

As shown in Figures 2A and B, FLT4 mRNA was not expressed in 7.5 day p.c. mouse embryos, but bright signals were detected in the posterior cardinal vein (cv) on day 8.5 of development. In contrast, the developing heart (data not shown) and dorsal aorta (da) were FLT4-negative. In the extraembryonic tissues FLT4 was prominently expressed in the allantois (al

in panel B), whereas developing blood islands of the yolk sac were negative (data not shown). On the other hand, angioblasts (ab) of the head mesenchyme were strongly FLT4 positive (C). In the developing placenta FLT4 signal was first seen in peripheral sinusoidal veins (data not shown). In  
5 9.5 day p.c. placenta the endothelium of venous lacunae (vl in D) and the giant cells partially fused to the Reichert's membrane (data not shown) expressed FLT4 mRNA.

Thus, although FLT4 expression was very prominent in the earliest endothelial cell precursors, the angioblasts, it appeared to be restricted only  
10 to certain vessels of 8.5 day p.c. embryos. The Tie receptor is known to be expressed in all endothelial cells of developing mouse embryos and thus provides a marker for these cells. Notably, in contrast to the Tie probe, the FLT4 probe hybridized very weakly if at all with arterial endothelia of 11.5 day p.c. embryos, e.g. with the endothelium of the developing dorsal aorta (da in  
15 Fig. 2 E,F) or the carotic arteries (data not shown). Instead, FLT4 signal was much more prominent in the developing veins. For example, FLT4 signal was detected veins surrounding the developing metanephros (v, sv in E), while the Tie probe predominantly recognized capillaries (c) within the metanephros (F).

20 As can be seen from Figure 3, FLT4 mRNA is distributed in several regions of a 12.5 day p.c. mouse embryo, being particularly prominent in the dilated vessel of the axillar region (ax). A similar FLT4 positive vessel structure was seen in the mid-sagittal section in the jugular area (data not shown). A plexus-like pattern of FLT4 expressing vessels appeared in the  
25 periorbital region (po) and surrounding the developing vertebrae (vb). Also, just beneath the developing skin, a FLT4-positive vascular network was evident (sc). Weaker capillary signals were obtained from several regions, including the developing brain (b). FLT4 mRNA could also be detected in small vessels of the neck region, of the developing snout and at the base of  
30 the developing tongue as well as in the tail region (data not shown). Besides, the liver (li) was strongly positive for FLT4 mRNA in a spotlike pattern.

During further development, FLT4 RNA appeared to become more restricted to certain vessels of the embryo. A 14.5 day p.c. embryo shows

nicely this restricted pattern of expression (Fig. 4 A,B). In the midsagittal section of Figure 4 the most prominent FLT4 signal is seen along the developing vertebral column in its anterior part. This signal was considered to originate from endothelial cells of the thoracic duct (dt), which is the largest lymphatic vessel formed at this time of development. In contrast, the dorsal aorta (da) and inferior vena cava (vc) were negative. Dilated vessels in the mesenteric region were also strongly positive for FLT4. Furthermore, as in the 12.5 day p.c. embryos, vessel networks along anatomical boundaries in the periorbital (po), lower jaw (lj) as well as in the neck (ne) region contained FLT4 positive endothelia. Similar structures were present in the pericardial space and throughout the subcutaneous (sc) tissue. Notably, in contrast to FLT4 negative vessels, all FLT4 positive vessels were devoid of blood cells in their lumen. These expression patterns suggest that FLT4 becomes confined to the endothelia of lymphatic vessels at this time of development. An additional site where we observed FLT4 expression, was in the sinusoids of the developing bone marrow (bm).

Photographs of a transverse section of the upper thorax of a 16.5 day p.c. embryo hybridized with the FLT4 probe are shown in panels C and D of Figure 4. The section shown in C has been stained with hematoxylin-eosin to visualize the different types of vessels in this area. These include the carotic and brachiocephalic arteries (ca, ba), the vena cava (vc) and the thoracic duct, which is smaller in size and lacks surrounding muscular and connective tissue (arrow). A magnification of the region of thoracic duct is shown in panel D, where the FLT4 autoradiographic grains can be seen. Endothelial cells of the thoracic duct as well as a small vessel (v) in the vicinity hybridize with the FLT4 probe.

#### EXAMPLE 4

##### Analysis of FLT4 mRNA in cultured endothelial cells

The *in situ* hybridization results described in Example 3, showed that FLT4 is expressed in venous endothelial cells and later in lymphatic vessels and some venous endothelial cells, but not in arterial endothelia. In order to see if such regulation was maintained *in vitro*, we studied cultured endothelial cells using Northern blotting and hybridization analysis.



Endothelial cells from human aorta, femoral vein, umbilical vein, and from foreskin microvessels were isolated, cultured and characterized as previously described by Van Hinsberg, <sup>(15, 16)</sup>. They were used at confluent density after five to eight passages (split ratio 1:3) for the isolation of polyadenylated RNA.

The endothelial cell lines EA-hy926, BCE and LEII did not express FLT4 (data not shown). However, cultured human microvascular, venous and umbilical vein endothelial cells were positive for the FLT4-specific 5.8 and 4.5 kb mRNAs, whereas the aortic endothelial cells were negative (Fig. 5). In contrast, another endothelial receptor tyrosine kinase gene, Tie, was expressed as a 4.4 kb mRNA in all endothelial cell types studied.

#### EXAMPLE 5

##### FLT4 mRNA in in adult human tissues

The results obtained in Example 3 indicated that the FLT4 mRNA becomes largely confined to the endothelium of lymphatic vessels during development. Because of the potential significance of this finding in humans, we also studied FLT4 in adult human tissues using a human FLT4 probe. The human FLT4 probe used was an EcoRI-SphI fragment covering base pairs 1-595 of the cDNA <sup>(3)</sup>. The von Willebrand factor probe was an EcoRI-HindIII fragment covering base pairs 1-2334 <sup>(17)</sup>.

We used routinely fixed material sent for histopathological diagnosis. Normal lung tissue was obtained from a resection of the left inferior lung lobe affected by epidermoid cancer. Mesenterium and mesenterial lymph nodes were obtained from a patient having a colonic adenocarcinoma. A normal lymph node adjacent to the salivary gland was enucleated because of its abnormal size. The tonsils from two patients and the two appendixes had no diagnostic changes. Two lymphangiomyomas and three cystic lymphangiomas were studied with similar results.

For human tissues, which were routine samples fixed with 10% formalin for histopathological diagnosis, the normal *in situ* protocol gave just background, whereas microwave treatment instead of proteinaase K enabled specific hybridization <sup>(18, 19)</sup>.

In the mesenterium, lung and appendix lymphatic endothelia (lv) gave FLT4 signals, while veins (v), arteries (a) and capillaries (c) were negative (Fig. 6A-D and data not shown). To study whether FLT4 is expressed in the HEVs, the tonsils were studied. Indeed, in the tonsils, FLT4 specific  
5 autoradiographic grains were detected in some HEVs (E, F).

#### EXAMPLE 6

##### Analysis of FLT4 mRNA in normal and metastatic lymph node and in lymphangioma

A portion of a human mesenterial lymph node (see Example 5) was  
10 analysed for FLT4 expression. The results are shown in Figure 7.

FLT4 is expressed in the lymphatic sinuses (ls) and afferent and efferent lymphatic vessels (data not shown). The same pattern is seen in a lymph node containing adenocarcinoma metastases (C, D). Some HEVs in both normal and metastatic lymph node were also positive. In panel E, FLT4  
15 expression is shown in a cystic lymphangioma (compare with the hematoxylin-eosin stained section in F). Notably, the specificity of FLT4 to lymphatic endothelia is evident from the comparison with the *in situ* signals for von Willebrandt factor in all blood vessels (F).

#### EXAMPLE 7

##### Localization of FLT4 in fetal endothelial cells

An FLT4 cDNA fragment encoding the 40 carboxy terminal amino acids of the short form was cloned as a 657 bp *EcoRI*-fragment into the pGEX-1IT bacterial expression vector (Pharmacia) in frame with the glutathione-S-transferase coding region. The resulting GST-FLT4 fusion  
25 protein was produced in *E.coli* and purified by affinity chromatography using a glutathione-Sepharose 4B column. The purified protein was lyophilized, dissolved in PBS, mixed with Freund's adjuvant and used for immunization of rabbits. Antisera were used after the fourth booster immunization.

Tissues from 17 and 20-week-old human fetuses were obtained from  
30 legal abortions induced with prostaglandins. The study was approved by the Ethical Committee of the Helsinki University Central Hospital. The

gestational age was estimated from the fetal foot length. The fetal tissues were embedded in Tissue-Tek (Miles), frozen immediately and stored at -70 °C.

Anti-FLT4 antiserum was cross-absorbed to a GST-Sepharose column to remove anti-GST-antibodies and then purified by GST-FLT4 affinity chromatography. Several 6 µm-thick cryostat sections of the tissues were fixed with acetone and treated with 0.3% H<sub>2</sub>O<sub>2</sub> in methanol for 30 min to block endogenous peroxidase activity. After washing, the sections were incubated with 5% normal swine serum. Sections were then incubated with antibodies against FLT4, washed and bound antibodies were detected with peroxidase-conjugated swine anti-rabbit IgG followed by staining for peroxidase activity using 0.2% 3,3-diaminobenzidine (Amersham) as a substrate. The sections were counterstained in Meyer's hematoxylin.

Anti-FLT4 immunoperoxidase staining of human fetal mesenterium showed FLT4 protein in the endothelium of several vessels (Fig. 4A), while control stainings with antigen-blocked anti-FLT4 antibodies (B) and preimmune sera (C) were negative. For comparison, Figure 4D shows results of staining using an antiserum against the factor VIII-related antigen, which is specific for vascular endothelial cells.

## EXAMPLE 8

### Production of monoclonal antibodies against FLT4

#### Fusion I:

Four months old Balb/c male mice were immunized by intraperitoneal injection of the recombinantly produced FLT4 protein (see Example 7) in concentrated medium (150 µg/mouse), emulsified with Freund's complete adjuvant. Booster injections of 150 µg were given at three to four week intervals and a final booster (10 µg FLT4 in PBS administered intraperitoneally) was given after another three-week interval. Four days after the final booster dose, the mice were sacrificed and mouse splenic lymphoid cells were fused with SP 2/0 plasmacytoma cells at a 2:1 ratio, respectively.

The fused cells were harvested in 96-well culture plates (NUNC) in Ex-Cell 320 medium (SERALAB) containing 20% fetal calf serum and HAT supplement (hypoxanthine-aminopterin-thymidine; GIBCO, 043-01060H; diluted 50-fold). Cells were cultured at +37°C, in a 5% CO<sub>2</sub> atmosphere.

- 5 After 10 days, HAT-supplemented medium was changed to HT-supplemented cell culture medium (GIBCO; 043-01065H, diluted 50-fold). HT medium is identical to HAT medium, but lacks aminopterin.

In three weeks specific antibody production was determined by the antigen-specific immunofluorometric assay, IFMA, described in Example 10

- 10 The master clones were cloned by limited dilutions as described by Staszewski et al., Yale Journal of Biology and Medicine, 57:865-868 (1984). Positive clones were expanded onto 24-well tissue culture plates (NUNC), recloned, and re-tested by the same method. Positive clones were tested by fluorescence-activated cell sorting (FACS).

- 15 The stable clones secreted immunoglobulins belonging to the IgG1 class, except one, which produced Ig probably belonging to class IgA. The subclass of monoclonal antibody was determined using rat monoclonal antibody to mouse subclass as biotin conjugate (SEROTEC) in IFMA.

- 20 Balb/c mice were used to produce monoclonal antibodies in ascites fluid. The hybridomas described above were intraperitoneally injected into mice after pretreatment of the animals with pristane (2,6,10,14-tetramethylpentadecan 98%, ALDRICH-CHEMIE D7924 Steinheim, Cat.No. T 2,280-2). 0.5 ml of pristane (i.v.) was injected about two weeks prior to the hybridoma cells. The amount of cells injected were approximately 7.5 to 9 x 10<sup>6</sup> per mouse. Ascites was collected 10 to 14 days after injection of the hybridomas.

#### Fusion II:

- 30 Two months old Balb/c mice (female) were immunized by intraperitoneal injection of the recombinantly produced FLT-4 protein (see Example 7) (20 µg/mouse), emulsified with Freund's complete adjuvant. Booster injections of 20 µg were given at three to four week intervals and a final booster (10 µg FLT-4 in PBS administered i.v.) was given after another three-week interval. Four days after the final booster dose, the mice were

sacrificed and mouse splenic lymphoid cells were fused with SP 2/0 plasmacytoma cells at a 2:1 ratio, respectively.

The fused cells were harvested in 96-well culture plates (FALCON) in OptiMEM 1 (with Glutamax 1, 51985-026, GIBCO BRL) medium containing  
5 20 % fetal calf serum and HAT supplement (hypoxanthine-aminopterin-thymidine; GIBCO BRL 21060-017; diluted 1:50 fold). Cells were cultured at +37 °C, in a 5% CO<sub>2</sub> atmosphere. After 10 days, HAT-supplemented medium was changed to HT-supplemented cell culture medium (GIBCO BRL; 41065-012, diluted 1:50-fold). HT-medium is identical to HAT-medium, but  
10 lacks aminopterin.

In three weeks specific antibody production was determined by the antigen-specific ImmunoFluoroMetric Assay (IFMA) described in Example 9. The master clones were cloned by limited dilutions as described by Staszewski et al., Yale Journal of Biology and Medicine, 57:865-868 (1984).  
15 Positive clones were expanded onto 24-well tissue culture plates (FALCON), recloned, and re-tested by the same method. Positive clones were tested by fluorescence-activated cell sorting (FACS).

The 2E11 and 6B2 clones secreted immunoglobulins belonging to the IgG<sub>1</sub> class, 2B12 clones produced Ig belonging to subclass IgM. The mouse  
20 subclass IgG<sub>1</sub> was determined using rat monoclonal antibody against mouse subclass heavy chain as biotin conjugate (SEROTEC) in IFMA and the mouse subclass IgM was determined with Mouse Monoclonal Antibody Isotyping Kit (Dipstick Format) (19663-012, Life Technologies Inc.).

#### EXAMPLE 9

##### 25 Specificity of monoclonal antibodies against FLT4

##### Fusion I antibodies:

The extracellular domain of FLT4 described in Example 7, was labelled according to Mukkala et al., in *Anal.Biochem.* 176(2):319-325, 1989, with the following modification: a 250 times molar excess of isothiocyanate  
30 DTTA-Eu (N1 chelate, WALLAC, Finland) was added to the FLT4 solution (0.5 mg/ml in PBS) and the pH was adjusted to about 9 by adding 0.5 mol/L

## 20

sodium carbonate buffer, pH 9.8. The labelling was performed overnight at +4°C. Unbound label was removed using PD-10 (PHARMACIA, Sweden) with TSA buffer (50 mmol/L Tris-HCl, pH 7.8 containing 0.15 mol/l NaCl) as eluent.

5           After purification, 1 mg/ml bovine serum albumin (BSA) was added to the labelled FLT4 and the label was stored at +4°C. The number of europium ions incorporated per FLT4 molecule was 1.9, as determined by measuring the fluorescence in a ratio to that of known EuCl<sub>3</sub> standards (Hemmilä et al., *Anal.Biochem.*, 137:335-343, 1984).

10           The antibodies produced in Example 8, were screened using a Sandwich type immunofluorometric assay using microtitration strip wells (NUNC, polysorb) coated with rabbit anti-mouse Ig (Z 259, DAKOPATTS). The precoated wells were washed once by Platelash 1296-024 (WALLAC) with DELFIA wash solution. The DELFIA assay buffer was used as a dilution  
15           buffer for cell culture supernatants and for serum of the spleenectomized mouse (at dilutions between 1:1000 to 1: 100 000) used as positive control in the preliminary screening assay.

          An overnight incubation at +4°C (or alternatively for 2 hours at room temperature) was begun by shaking on a Plateshake shaker (1296-001,  
20           WALLAC) for 5 min followed by washing four times with wash solution as described above.

          The europium-labelled FLT4 was added at a dilution of 1:500 in 100 µl of the assay buffer. After 5 min on a Plateshake shaker and one hour incubation at RT the strips were washed as described above.

25           Enhancement solution (DELFIA) was added at 200 µl/well. The plates were then shaken for 5 min on a Plateshake shaker and the intensity of fluorescence was measured by ARCUS-1230 (WALLAC) for 10-15 min. (Lövgren et al., In: Collins W.P. (Ed) *Alternative Immunoassays*, John Wiley & Sons Ltd, 1985; pp. 203-216).

30           The resulting monoclonal antibodies against FLT4 and corresponding FACS results are summarized in Table 2.

**TABLE 2**

Mab clones	LTR% <sup>a)</sup>	NEO% <sup>b)</sup>	DELFIA-counts
1B1	67,3	1	20625
1B1D11	75	1,2	19694
1B1F8	76,1	1,4	18580
4F6	69,9	1,2	23229
4F6B8G12	75	0,3	24374
4F6B8H11	75,9	0,3	28281
4F6B8E12	74,8	0,4	27097
4F6B8G10	75,3	0,4	26063
9D9	45,1	0,75	17316
9D9D10	71,7	2,3	18230
9D9F9	73	1,8	11904
9D9G6	74,3	2,9	16743
9D9G7	70,7	1,3	17009
10E4	24,2	1,4	39202
10E4B10E12	32,3	0,3	42490
10E4B10G10	36,5	0,3	54815
10E4B10F12	45,6	0,4	43909
10E4B10G12	45,7	0,5	35576
11G2	30,2	1,6	11304
11G2D12	74,4	1,5	14660
11G2G9	74,2	0,9	10283
11G2H7	74,4	2,1	25382

a) FACS results with LTR transfected cells

b) FACS results with NEO cells (control)

- 5 One clone, designated anti-FLT4 9D9F9 was found to stably secrete monoclonal antibody which was determined to be of immunoglobulin class IgG1 by IFMA. Hybridoma 9d9f9 was deposited with the German Collection of Microorganisms and Cell Cultures, Department of Human and Animal Cell Cultures and Viruses, Mascheroder Weg 1b, 3300 Braunschweig, Germany,
- 10 March 23, 1995, and given accession No. ACC2210.

Fusion II antibodies:

The extracellular domain of FLT-4 described in Example 7, was labelled according to Mukkala et al., in Anal.Biochem. 176(2): 319-325, 1989, with the following modification: a 250 times molar excess of isothiocyanate  
5 DTTA-Eu (N1 chelate, Wallac, Finland) was added to the FLT-4 solution (0.5 mg/ml in PBS) and the pH was adjusted to about 9 by adding 0.5 mol/L sodium carbonate buffer, pH 9.8. The labelling was performed overnight at +4°C. Unbound label was removed using PD-10 (PHARMACIA) with TSA buffer (50 mmol/L Tris-HCl, pH 7.8 containing 0.15 mol/L NaCl) as eluent.

10 After purification, 1 mg/ml bovine serum albumin (BSA) was added to the labelled FLT-4 and the label was stored at +4°C. The number of europium ions incorporated per FLT-4 molecule was 1.9, as determined by measuring the fluorescence in a ratio to that of known EuCl<sub>3</sub> standards (Hemmil et al., Anal.Biochem., 137: 335-343, 1984).

15 The antibodies produced in Example 8, were screened using a FLT-4 specific IFMA using microtitration wells (Nunc, Polysorb) coated with rabbit antimouse Ig (Z 259, DAKO). The precoated wells were washed once with wash solution (Wallac) by using DELFIA Plate wash.

The DELFIA assay buffer was used as dilution buffer for cell culture  
20 supernatants (dilution 1:2 in preliminary screening) and for serum of the splenectomized mouse (dilutions 1:1 000 to 1:100 000) which was used as positive control. As standard the purified antiFLT-4 9D9F9 (mouse subclass IgG1 ) was used at concentrations between 1.0 ng/ml and 250 ng/ml. Samples were first shaken at room temperature for five minutes on Plate  
25 shake (Wallac) and then incubated approx. 18 hours at +4°C. The frames were first washed four times, then the Eu-labelled FLT-4 (1:2000, in 100 µl assay buffer) was added and finally the frames were incubated for one hour at room temperature. After washing as described the enhancement solution (200 µl/well, Wallac) was added and the frames were shaken for 5 minutes  
30 on Plate shake. The intensity of fluorescence was measured by ARCUS-1230 (Wallac).



## 23

The resulting monoclonal antibodies against FLT-4 and corresponding results are summarized in Table 3.

A standard curve for quantitation of antiFLT-4 antibodies was made by using affinity purified antiFLT-4 9D9F9. The linear range reached from 1.0 ng/ml to 250 ng/ml.

Cell lysate of NIH 3T3 cells cotransfected with pLTRFLT4 construct expressing full-length FLT4 on the surface was electrophoresed in 6.5% SDS-PAGE, proteins were transferred onto nitrocellulose nitrate membrane (0.45  $\mu$ m, SCHLEICHER & SCHUELL) and immunoblotted with Mab cell culture supernatants (1:10, 50 mmol/L TRIS - 40 mmol/L glycine buffer containing methanol 4%, SDS 0.04%). The specificity of Mab was detected using incubation with HRP-conjugated rabbit antimouse Ig (P 161, DAKO, diluted 1:1000 in 20 mmol/L TRIS buffer pH 7.5 containing 150 mmol/L saline, 5% milk powder) and ECL (Enhanced chemiluminescence, AMERSHAM).

TABLE 3.

	Mab clones	LTR %a)	NEOb)	approx.Mab prod. ng/ml/106 cells	WB
20	2B12E10	39.5	6.0	440	+
	2E11D11	44.6	8.8	110	+
	2E11F9	49.5	4.5	100	+
	2E11F12	46.0	4.1	180	+
25	2E11G8	41.2	7.8	160	+
	6B2E12	NF	NF	1390	+
	6B2F8	NF	NF	470	+
	6B2G6	NF	NF	630	+
30	6B2H5	NF	NF	740	+
	6B2H8	NF	NF	1800	+

a) FACS results with LTR transfected cells

b) FACS results with NEO cells (control)

35 NF not functioning in FACS

c) quantitation of Mab production based on affinity purified antiFLT 9D9F9 antibody used as standard

As is evident from the foregoing, antibodies according to the present invention are useful in the diagnosis and identification of lymphatic vessels, lymphatic endothelial cells, high endothelial venules, lymphangiomas, metastatic lymph nodes and other disease states of the lymphatic system, the  
5 detection and monitoring of metastatic spread, in the stimulation and inhibition of endothelial cells of lymphatic vessels and high endothelial venules, in the introduction of molecules selectively into endothelial cells and in the imaging of lymphatic vessels and their disease states. Other uses of the presently-claimed subject matter are apparent to the skilled artisan.

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A. The indications made below relate to the microorganism referred to in the description</b> on page <u>3</u> , line <u>13 - 17</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution DSM-DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELLKULTUREN GmbH	
Address of depositary institution (including postal code and country) Mascheroder Weg 1b D-38124 Braunschweig Germany	
Date of deposit 1995-03-23	Accession Number DSM ACC2210
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
As regards the respective Patent Offices of the respective designated states, the applicant requests that a sample of the deposited microorganisms only be made available to an expert nominated by the requester until the date on which the patent is granted or the date on which the application has been refused or withdrawn or is deemed to be withdrawn	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="border-bottom: 1px solid black; padding-bottom: 5px;"> <b>For receiving Office use only</b> </div> <div style="padding: 5px;"> <input checked="" type="checkbox"/> This sheet was received with the international application         </div> <div style="border-top: 1px solid black; padding-top: 5px;">         Authorized officer  </div>	<div style="border-bottom: 1px solid black; padding-bottom: 5px;"> <b>For International Bureau use only</b> </div> <div style="padding: 5px;"> <input type="checkbox"/> This sheet was received by the International Bureau on:         </div> <div style="border-top: 1px solid black; padding-top: 5px;">         Authorized officer       </div>
--	---

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: Alitalo, Kari  
(B) STREET: Nyyrikintie 4 A  
(C) CITY: ESPOO  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-02100

(A) NAME: Kaipainen, Arja  
(B) STREET: Messeniuksenkatu 7 A 9  
(C) CITY: HELSINKI  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-00250

(A) NAME: Korhonen, Jaana  
(B) STREET: Agricolankatu 7 C 68  
(C) CITY: HELSINKI  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-00530

(A) NAME: Mustonen, Tuija  
(B) STREET: Pihlajatie 8 A 2  
(C) CITY: HELSINKI  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-00270

(A) NAME: Pajusola, Katri  
(B) STREET: Kasteholmantie 4 A 8  
(C) CITY: HELSINKI  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-00900

(A) NAME: Matikainen, Marja-Terttu  
(B) STREET: Lankkistentanhua 12  
(C) CITY: MYNÄMÄKI  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-23100

(A) NAME: Karnani, Päivi  
(B) STREET: Mullintie 10 B 62  
(C) CITY: TURKU  
(E) COUNTRY: FINLAND  
(F) POSTAL CODE (ZIP): FIN-20300

(ii) TITLE OF INVENTION: FLT4 RECEPTOR TYROSINE KINASE AND ITS USE  
IN DIAGNOSIS AND THERAPY

(iii) NUMBER OF SEQUENCES: 1

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: WO TO BE ASSIGNED

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/257754

(B) FILING DATE: 9-JUN-1994

(2) INFORMATION FOR SEQ ID NO: 1:

CCACGCGCAG CGGCCGGAG ATG CAG CGG GGC GCC GCG CTG TGC CTG CGA CTG  
52

Met Gln Arg Gly Ala Ala Leu Cys Leu Arg Leu

1

5

10

TGG CTC TGC CTG GGA CTC CTG GAC GGC CTG GTG AGT GGC TAC TCC ATG  
100

Trp Leu Cys Leu Gly Leu Leu Asp Gly Leu Val Ser Gly Tyr Ser Met  
15 20 25

ACC CCC CCG ACC TTG AAC ATC ACG GAG GAG TCA CAC GTC ATC GAC ACC  
148

Thr Pro Pro Thr Leu Asn Ile Thr Glu Glu Ser His Val Ile Asp Thr  
30 35 40

GGT GAC AGC CTG TCC ATC TCC TGC AGG GGA CAG CAC CCC CTC GAG TGG  
196

Gly Asp Ser Leu Ser Ile Ser Cys Arg Gly Gln His Pro Leu Glu Trp  
45 50 55

GCT TGG CCA GGA GCT CAG GAG GCG CCA GCC ACC GGA GAC AAG GAC AGC  
244

Ala Trp Pro Gly Ala Gln Glu Ala Pro Ala Thr Gly Asp Lys Asp Ser  
60 65 70 75

GAG GAC ACG GGG GTG GTG CGA GAC TGC GAG GGC ACA GAC GCC AGG CCC  
292

Glu Asp Thr Gly Val Val Arg Asp Cys Glu Gly Thr Asp Ala Arg Pro  
80 85 90

TAC TGC AAG GTG TTG CTG CTG CAC GAG GTA CAT GCC AAC GAC ACA GGC  
340

Tyr Cys Lys Val Leu Leu Leu His Glu Val His Ala Asn Asp Thr Gly  
95 100 105

AGC TAC GTC TGC TAC TAC AAG TAC ATC AAG GCA CGC ATC GAG GGC ACC  
388

Ser Tyr Val Cys Tyr Tyr Lys Tyr Ile Lys Ala Arg Ile Glu Gly Thr  
110 115 120

ACG GCC GCC AGC TCC TAC GTG TTC GTG AGA GAC TTT GAG CAG CCA TTC  
436

Thr Ala Ala Ser Ser Tyr Val Phe Val Arg Asp Phe Glu Gln Pro Phe  
125 130 135

ATC AAC AAG CCT GAC ACG CTC TTG GTC AAC AGG AAG GAC GCC ATG TGG  
484

Ile Asn Lys Pro Asp Thr Leu Leu Val Asn Arg Lys Asp Ala Met Trp  
140 145 150 155

GTG CCC TGT CTG GTG TCC ATC CCC GGC CTC AAT GTC ACG CTG CGC TCG  
532

Val Pro Cys Leu Val Ser Ile Pro Gly Leu Asn Val Thr Leu Arg Ser  
160 165 170

CAA AGC TCG GTG CTG TGG CCA GAC GGG CAG GAG GTG GTG TGG GAT GAC  
580

Gln Ser Ser Val Leu Trp Pro Asp Gly Gln Glu Val Val Trp Asp Asp  
175 180 185

CGG CGG GGC ATG CTC GTG TCC ACG CCA CTG CTG CAC GAT GCC CTG TAC  
 628  
 Arg Arg Gly Met Leu Val Ser Thr Pro Leu Leu His Asp Ala Leu Tyr  
 190 195 200

CTG CAG TGC GAG ACC ACC TGG GGA GAC CAG GAC TTC CTT TCC AAC CCC  
 676  
 Leu Gln Cys Glu Thr Thr Trp Gly Asp Gln Asp Phe Leu Ser Asn Pro  
 205 210 215

TTC CTG GTG CAC ATC ACA GGC AAC GAG CTC TAT GAC ATC CAG CTG TTG  
 724  
 Phe Leu Val His Ile Thr Gly Asn Glu Leu Tyr Asp Ile Gln Leu Leu  
 220 225 230 235

CCC AGG AAG TCG CTG GAG CTG CTG GTA GGG GAG AAG CTG GTC CTG AAC  
 772  
 Pro Arg Lys Ser Leu Glu Leu Leu Val Gly Glu Lys Leu Val Leu Asn  
 240 245 250

TGC ACC GTG TGG GCT GAG TTT AAC TCA GGT GTC ACC TTT GAC TGG GAC  
 820  
 Cys Thr Val Trp Ala Glu Phe Asn Ser Gly Val Thr Phe Asp Trp Asp  
 255 260 265

TAC CCA GGG AAG CAG GCA GAG CGG GGT AAG TGG GTG CCC GAG CGA CGC  
 868  
 Tyr Pro Gly Lys Gln Ala Glu Arg Gly Lys Trp Val Pro Glu Arg Arg  
 270 275 280

TCC CAG CAG ACC CAC ACA GAA CTC TCC AGC ATC CTG ACC ATC CAC AAC  
 916  
 Ser Gln Gln Thr His Thr Glu Leu Ser Ser Ile Leu Thr Ile His Asn  
 285 290 295

GTC AGC CAG CAC GAC CTG GGC TCG TAT GTG TGC AAG GCC AAC AAC GGC  
 964  
 Val Ser Gln His Asp Leu Gly Ser Tyr Val Cys Lys Ala Asn Asn Gly  
 300 305 310 315

ATC CAG CGA TTT CGG GAG AGC ACC GAG GTC ATT GTG CAT GAA AAT CCC  
 1012  
 Ile Gln Arg Phe Arg Glu Ser Thr Glu Val Ile Val His Glu Asn Pro  
 320 325 330

TTC ATC AGC GTC GAG TGG CTC AAA GGA CCC ATC CTG GAG GCC ACG GCA  
 1060  
 Phe Ile Ser Val Glu Trp Leu Lys Gly Pro Ile Leu Glu Ala Thr Ala  
 335 340 345

GGA GAC GAG CTG GTG AAG CTG CCC GTG AAG CTG GCA GCG TAC CCC CCG  
 1108  
 Gly Asp Glu Leu Val Lys Leu Pro Val Lys Leu Ala Ala Tyr Pro Pro  
 350 355 360

CCC GAG TTC CAG TGG TAC AAG GAT GGA AAG GCA CTG TCC GGG CGC CAC  
 1156  
 Pro Glu Phe Gln Trp Tyr Lys Asp Gly Lys Ala Leu Ser Gly Arg His  
 365 370 375

AGT CCA CAT GCC CTG GTG CTC AAG GAG GTG ACA GAG GCC AGC ACA GGC  
 1204  
 Ser Pro His Ala Leu Val Leu Lys Glu Val Thr Glu Ala Ser Thr Gly  
 380 385 390 395

ACC TAC ACC CTC GCC CTG TGG AAC TCC GCT GCT GGC CTG AGG CGC AAC  
 1252  
 Thr Tyr Thr Leu Ala Leu Trp Asn Ser Ala Ala Gly Leu Arg Arg Asn  
 400 405 410

ATC AGC CTG GAG CTG GTG GTG AAT GTG CCC CCC CAG ATA CAT GAG AAG  
 1300  
 Ile Ser Leu Glu Leu Val Val Asn Val Pro Pro Gln Ile His Glu Lys  
 415 420 425

GAG GCC TCC TCC CCC AGC ATC TAC TCG CGT CAC AGC CGC CAG GCC CTC  
 1348  
 Glu Ala Ser Ser Pro Ser Ile Tyr Ser Arg His Ser Arg Gln Ala Leu  
 430 435 440

ACC TGC ACG GCC TAC GGG GTG CCC CTG CCT CTC AGC ATC CAG TGG CAC  
 1396  
 Thr Cys Thr Ala Tyr Gly Val Pro Leu Pro Leu Ser Ile Gln Trp His  
 445 450 455

TGG CGG CCC TGG ACA CCC TGC AAG ATG TTT GCC CAG CGT AGT CTC CGG  
 1444

Trp Arg Pro Trp Thr Pro Cys Lys Met Phe Ala Gln Arg Ser Leu Arg  
 460 465 470 475

CGG CGG CAG CAG CAA GAC CTC ATG CCA CAG TGC CGT GAC TGG AGG GCG  
 1492  
 Arg Arg Gln Gln Gln Asp Leu Met Pro Gln Cys Arg Asp Trp Arg Ala  
 480 485 490

GTG ACC ACG CAG GAT GCC GTG AAC CCC ATC GAG AGC CTG GAC ACC TGG  
 1540  
 Val Thr Thr Gln Asp Ala Val Asn Pro Ile Glu Ser Leu Asp Thr Trp  
 495 500 505

ACC GAG TTT GTG GAG GGA AAG AAT AAG ACT GTG AGC AAG CTG GTG ATC  
 1588  
 Thr Glu Phe Val Glu Gly Lys Asn Lys Thr Val Ser Lys Leu Val Ile  
 510 515 520

CAG AAT GCC AAC GTG TCT GCC ATG TAC AAG TGT GTG GTC TCC AAC AAG  
 1636  
 Gln Asn Ala Asn Val Ser Ala Met Tyr Lys Cys Val Val Ser Asn Lys  
 525 530 535

GTG GGC CAG GAT GAG CGG CTC ATC TAC TTC TAT GTG ACC ACC ATC CCC  
 1684  
 Val Gly Gln Asp Glu Arg Leu Ile Tyr Phe Tyr Val Thr Thr Ile Pro  
 540 545 550 555

GAC GGC TTC ACC ATC GAA TCC AAG CCA TCC GAG GAG CTA CTA GAG GGC  
 1732  
 Asp Gly Phe Thr Ile Glu Ser Lys Pro Ser Glu Glu Leu Leu Glu Gly  
 560 565 570

CAG CCG GTG CTC CTG AGC TGC CAA GCC GAC AGC TAC AAG TAC GAG CAT  
 1780  
 Gln Pro Val Leu Leu Ser Cys Gln Ala Asp Ser Tyr Lys Tyr Glu His  
 575 580 585

CTG CGC TGG TAC CGC CTC AAC CTG TCC ACG CTG CAC GAT GCG CAC GGG  
 1828  
 Leu Arg Trp Tyr Arg Leu Asn Leu Ser Thr Leu His Asp Ala His Gly  
 590 595 600

AAC CCG CTT CTG CTC GAC TGC AAG AAC GTG CAT CTG TTC GCC ACC CCT  
 1876  
 Asn Pro Leu Leu Leu Asp Cys Lys Asn Val His Leu Phe Ala Thr Pro  
 605 610 615

CTG GCC GCC AGC CTG GAG GAG GTG GCA CCT GGG GCG CGC CAC GCC ACG  
 1924  
 Leu Ala Ala Ser Leu Glu Glu Val Ala Pro Gly Ala Arg His Ala Thr  
 620 625 630 635

CTC AGC CTG AGT ATC CCC CGC GTC GCG CCC GAG CAC GAG GGC CAC TAT  
 1972  
 Leu Ser Leu Ser Ile Pro Arg Val Ala Pro Glu His Glu Gly His Tyr  
 640 645 650

GTG TGC GAA GTG CAA GAC CGG CGC AGC CAT GAC AAG CAC TGC CAC AAG  
 2020  
 Val Cys Glu Val Gln Asp Arg Arg Ser His Asp Lys His Cys His Lys  
 655 660 665

AAG TAC CTG TCG GTG CAG GCC CTG GAA GCC CCT CGG CTC ACG CAG AAC  
 2068  
 Lys Tyr Leu Ser Val Gln Ala Leu Glu Ala Pro Arg Leu Thr Gln Asn  
 670 675 680

TTG ACC GAC CTC CTG GTG AAC GTG AGC GAC TCG CTG GAG ATG CAG TGC  
 2116  
 Leu Thr Asp Leu Leu Val Asn Val Ser Asp Ser Leu Glu Met Gln Cys  
 685 690 695

TTG GTG GCC GGA GCG CAC GCG CCC AGC ATC GTG TGG TAC AAA GAC GAG  
 2164  
 Leu Val Ala Gly Ala His Ala Pro Ser Ile Val Trp Tyr Lys Asp Glu  
 700 705 710 715

AGG CTG CTG GAG GAA AAG TCT GGA GTC GAC TTG GCG GAC TCC AAC CAG  
 2212  
 Arg Leu Leu Glu Glu Lys Ser Gly Val Asp Leu Ala Asp Ser Asn Gln  
 720 725 730

AAG CTG AGC ATC CAG CGC GTG CGC GAG GAG GAT GCG GGA CGC TAT CTG  
 2260  
 Lys Leu Ser Ile Gln Arg Val Arg Glu Glu Asp Ala Gly Arg Tyr Leu  
 735 740 745

TGC AGC GTG TGC AAC GCC AAG GGC TGC GTC AAC TCC TCC GCC AGC GTG  
 2308  
 Cys Ser Val Cys Asn Ala Lys Gly Cys Val Asn Ser Ser Ala Ser Val  
 750 755 760

GCC GTG GAA GGC TCC GAG GAT AAG GGC AGC ATG GAG ATC GTG ATC CTT  
 2356  
 Ala Val Glu Gly Ser Glu Asp Lys Gly Ser Met Glu Ile Val Ile Leu  
 765 770 775

GTC GGT ACC GGC GTC ATC GCT GTC TTC TTC TGG GTC CTC CTC CTC CTC  
 2404  
 Val Gly Thr Gly Val Ile Ala Val Phe Phe Trp Val Leu Leu Leu Leu  
 780 785 790 795

ATC TTC TGT AAC ATG AGG AGG CCG GCC CAC GCA GAC ATC AAG ACG GGC  
 2452  
 Ile Phe Cys Asn Met Arg Arg Pro Ala His Ala Asp Ile Lys Thr Gly  
 800 805 810



TAC CTG TCC ATC ATC ATG GAC CCC GGG GAG GTG CCT CTG GAG GAG CAA  
 2500  
 Tyr Leu Ser Ile Ile Met Asp Pro Gly Glu Val Pro Leu Glu Glu Gln  
 815 820 825

TGC GAA TAC CTG TCC TAC GAT GCC AGC CAG TGG GAA TTC CCC CGA GAG  
 2548  
 Cys Glu Tyr Leu Ser Tyr Asp Ala Ser Gln Trp Glu Phe Pro Arg Glu  
 830 835 840

CGG CTG CAC CTG GGG AGA GTG CTC GGC TAC GGC GCC TTC GGG AAG GTG  
 2596  
 Arg Leu His Leu Gly Arg Val Leu Gly Tyr Gly Ala Phe Gly Lys Val  
 845 850 855

GTG GAA GCC TCC GCT TTC GGC ATC CAC AAG GGC AGC AGC TGT GAC ACC  
 2644  
 Val Glu Ala Ser Ala Phe Gly Ile His Lys Gly Ser Ser Cys Asp Thr  
 860 865 870 875

GTG GCC GTG AAA ATG CTG AAA GAG GGC GCC ACG GCC AGC GAG CAC CGC  
 2692  
 Val Ala Val Lys Met Leu Lys Glu Gly Ala Thr Ala Ser Glu His Arg  
 880 885 890

GCG CTG ATG TCG GAG CTC AAG ATC CTC ATT CAC ATC GGC AAC CAC CTC  
 2740  
 Ala Leu Met Ser Glu Leu Lys Ile Leu Ile His Ile Gly Asn His Leu  
 895 900 905

AAC GTG GTC AAC CTC CTC GGG GCG TGC ACC AAG CCG CAG GGC CCC CTC  
 2788  
 Asn Val Val Asn Leu Leu Gly Ala Cys Thr Lys Pro Gln Gly Pro Leu  
 910 915 920

ATG GTG ATC GTG GAG TTC TGC AAG TAC GGC AAC CTC TCC AAC TTC CTG  
 2836  
 Met Val Ile Val Glu Phe Cys Lys Tyr Gly Asn Leu Ser Asn Phe Leu  
 925 930 935

CGC GCC AAG CGG GAC GCC TTC AGC CCC TGC GCG GAG AAG TCT CCC GAG  
 2884  
 Arg Ala Lys Arg Asp Ala Phe Ser Pro Cys Ala Glu Lys Ser Pro Glu  
 940 945 950 955

CAG CGC GGA CGC TTC CGC GCC ATG GTG GAG CTC GCC AGG CTG GAT CGG  
 2932  
 Gln Arg Gly Arg Phe Arg Ala Met Val Glu Leu Ala Arg Leu Asp Arg  
 960 965 970

AGG CGG CCG GGG AGC AGC GAC AGG GTC CTC TTC GCG CGG TTC TCG AAG  
 2980  
 Arg Arg Pro Gly Ser Ser Asp Arg Val Leu Phe Ala Arg Phe Ser Lys  
 975 980 985

ACC GAG GGC GGA GCG AGG CGG GCT TCT CCA GAC CAA GAA GCT GAG GAC  
 3028  
 Thr Glu Gly Gly Ala Arg Arg Ala Ser Pro Asp Gln Glu Ala Glu Asp  
 990 995 1000

CTG TGG CTG AGC CCG CTG ACC ATG GAA GAT CTT GTC TGC TAC AGC TTC  
 3076  
 Leu Trp Leu Ser Pro Leu Thr Met Glu Asp Leu Val Cys Tyr Ser Phe  
 1005 1010 1015

CAG GTG GCC AGA GGG ATG GAG TTC CTG GCT TCC CGA AAG TGC ATC CAC  
 3124  
 Gln Val Ala Arg Gly Met Glu Phe Leu Ala Ser Arg Lys Cys Ile His  
 1020 1025 1030 1035

AGA GAC CTG GCT GCT CGG AAC ATT CTG CTG TCG GAA AGC GAC GTG GTG  
 3172  
 Arg Asp Leu Ala Ala Arg Asn Ile Leu Leu Ser Glu Ser Asp Val Val  
 1040 1045 1050

AAG ATC TGT GAC TTT GGC CTT GCC CGG GAC ATC TAC AAA GAC CCT GAC  
 3220  
 Lys Ile Cys Asp Phe Gly Leu Ala Arg Asp Ile Tyr Lys Asp Pro Asp  
 1055 1060 1065

TAC GTC CGC AAG GGC AGT GCC CGG CTG CCC CTG AAG TGG ATG GCC CCT  
 3268  
 Tyr Val Arg Lys Gly Ser Ala Arg Leu Pro Leu Lys Trp Met Ala Pro  
 1070 1075 1080

GAA AGC ATC TTC GAC AAG GTG TAC ACC ACG CAG AGT GAC GTG TGG TCC  
 3316  
 Glu Ser Ile Phe Asp Lys Val Tyr Thr Thr Gln Ser Asp Val Trp Ser  
 1085 1090 1095

TTT GGG GTG CTT CTC TGG GAG ATC TTC TCT CTG GGG GCC TCC CCG TAC  
 3364  
 Phe Gly Val Leu Leu Trp Glu Ile Phe Ser Leu Gly Ala Ser Pro Tyr  
 1100 1105 1110 1115

CCT GGG GTG CAG ATC AAT GAG GAG TTC TGC CAG CGG CTG AGA GAC GGC  
 3412  
 Pro Gly Val Gln Ile Asn Glu Glu Phe Cys Gln Arg Leu Arg Asp Gly  
 1120 1125 1130

ACA AGG ATG AGG GCC CCG GAG CTG GCC ACT CCC GCC ATA CGC CGC ATC  
 3460  
 Thr Arg Met Arg Ala Pro Glu Leu Ala Thr Pro Ala Ile Arg Arg Ile  
 1135 1140 1145

ATG CTG AAC TGC TGG TCC GGA GAC CCC AAG GCG AGA CCT GCA TTC TCG  
 3508  
 Met Leu Asn Cys Trp Ser Gly Asp Pro Lys Ala Arg Pro Ala Phe Ser  
 1150 1155 1160

GAG CTG GTG GAG ATC CTG GGG GAC CTG CTC CAG GGC AGG GCC CTG CAA  
 3556  
 Glu Leu Val Glu Ile Leu Gly Asp Leu Leu Gln Gly Arg Gly Leu Gln  
 1165 1170 1175

GAG GAA GAG GAG GTC TGC ATG GCC CCG CGC AGC TCT CAG AGC TCA GAA  
 3604  
 Glu Glu Glu Glu Val Cys Met Ala Pro Arg Ser Ser Gln Ser Ser Glu  
 1180 1185 1190 1195

GAG GGC AGC TTC TCG CAG GTG TCC ACC ATG GCC CTA CAC ATC GCC CAG  
 3652  
 Glu Gly Ser Phe Ser Gln Val Ser Thr Met Ala Leu His Ile Ala Gln  
 1200 1205 1210

GCT GAC GCT GAG GAC AGC CCG CCA AGC CTG CAG CGC CAC AGC CTG GCC  
 3700  
 Ala Asp Ala Glu Asp Ser Pro Pro Ser Leu Gln Arg His Ser Leu Ala  
 1215 1220 1225

GCC AGG TAT TAC AAC TGG GTG TCC TTT CCC GGG TGC CTG GCC AGA GGG  
3748

Ala Arg Tyr Tyr Asn Trp Val Ser Phe Pro Gly Cys Leu Ala Arg Gly  
1230 1235 1240

GCT GAG ACC CGT GGT TCC TCC AGG ATG AAG ACA TTT GAG GAA TTC CCC  
3796

Ala Glu Thr Arg Gly Ser Ser Arg Met Lys Thr Phe Glu Glu Phe Pro  
1245 1250 1255

ATG ACC CCA ACG ACC TAC AAA GGC TCT GTG GAC AAC CAG ACA GAC AGT  
3844

Met Thr Pro Thr Thr Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp Ser  
1260 1265 1270 1275

GGG ATG GTG CTG GCC TCG GAG GAG TTT GAG CAG ATA GAG AGC AGG CAT  
3892

Gly Met Val Leu Ala Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg His  
1280 1285 1290

AGA CAA GAA AGC GGC TTC AGG TAGCTGAAGC AGAGAGAGAG AAGGCAGCAT  
3943

Arg Gln Glu Ser Gly Phe Arg  
1295

ACGTCAGCAT TTTCTTCTCT GCACTTATAA GAAAGATCAA AGACTTTAAG ACTTTTCGCTA  
4003

TTTCTTCTAC TGCTATCTAC TACAACTTC AAAGAGGAAC CAGGAGGACA AGAGGAGCAT  
4063

GAAAGTGGAC AAGGAGTGTG ACCACTGAAG CACCACAGGG AAGGGGTTAG GCCTCCGGAT  
4123

GACTGCGGGC AGGCCTGGAT AATATCCAGC CTCCCACAAG AAGCTGGTGG AGCAGAGTGT  
4183

TCCCTGACTC CT  
4195

## REFERENCES

1. Sabin, F.R. 1909. The lymphatic system in human embryos, with consideration of the morphology of the system as a whole. *Am. J. Anat.* 9:43.
2. van der Putte, S.C.J. 1975. The development of the lymphatic system in man. *Adv. Anat. Embryol. Cell Biol.* 51:3.
3. Pajusola, K., O. Aprelikova, J. Korhonen, A. Kaipainen, L. Pertovaara, R. Alitalo, and K. Alitalo. 1992. FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines. *Cancer Res.* 52:5738.
4. Kaipainen, A., J. Korhonen, K. Pajusola, O. Aprelikova, M.G. Persico, B.I. Terman, and K. Alitalo. 1993. The Related FLT4, FLT1 and KDR receptor tyrosine kinases show distinct expression patterns in human fetal endothelial cells. *J. Exp. Med.* 178:2077.
5. Galland, F., A. Karamysheva, M.-J. Pebusque, J.-P. Borg, R. Rottapel, P. Dubreuil, O. Rosnet, and D. Birnbaum. 1993. The *FLT4* gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor. *Oncogene.* 8:1233.
6. Millauer, B., S. Witzmann-Voos, H. Schnürch, R. Martinez, N.-P.H. Moller, W. Risau, and A. Ullrich. 1993. High affinity VEGF binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell.* 72:835.
7. Yamaguchi, T.P., D. Dumont, R.A. Conlon, M.L. Breitman, and J. Rossant. 1993. *flk-1*, an *flt*-related tyrosine kinase is an early marker for endothelial cell precursors. *Development.* 118:489.
8. Peters, K.G., C. De Vries, and L.T. Williams. 1993. Vascular endothelial growth factor receptor expression during embryogenesis and tissue repair suggests a role in endothelial differentiation and blood vessel growth. *Proc. Natl. Acad. Sci. USA* 90:8915.

9. Finnerty, H., K. Kelleher, G. Morris E., K. Bean, D. Merberg, R. Kritz, J. Morris C., H. Sookdeo, K.J. Turner, and C.R. Wood 1993. Molecular cloning of murine FLT and FLT4. *Oncogene*. 8:2293.
10. Korhonen, J., A. Polvi, J. Partanen, and K. Alitalo. 1993. The mouse *tie* receptor tyrosine kinase gene: expression during embryonic angiogenesis. *Oncogene*. 8:395.
11. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156.
- 10 12. Tokunaga, K., H. Taniguchi, K. Yoda, M. Shimizu, and S. Sakiyama. 1986. Nucleotide sequence of a full-length cDNA for mouse cytoskeletal beta-actin mRNA. *Nucleic. Acid. Res.* 14:2829.
13. Wilkinson, D.G., J.A. Bailes, J.E. Champion, and A.P. McMahon. 1987. A molecular analysis of mouse development from 8 to 10 days post coitum  
15 detects changes only in embryonic globin expression. *Development*. 99:493.
14. Wilkinson, D.G., J.A. Bailes, and A.P. McMahon. 1987. Expression of the proto-oncogene int-1 is restricted to specific neural cells in the developing mouse embryo. *Cell*. 50:79.
15. Van Hinsberg, V.W.M., D. Binnema, M.A. Scheffer, E.D. Sprengers, T. Kooistra, and D.C. Rijken. 1987. Production of plasminogen activators and inhibitors by serially propagated endothelial cells from adult human blood vessels. *Arteriosclerosis*. 7:389.
- 20 16. Van Hinsberg, V.W.M., M.A. Scheffer, and T. Kooistra. 1987. Effect of thrombin on the production of plasminogen activators and PA inhibitor-1 by human foreskin microvascular endothelial cells. *Thromb. Haemostas.* 57:148.
- 25 17. Bonthron, D.T., E.C. Orr, L.M. Mitsock, D. Ginsberg, R.I. Handin, and S.H. Orkin. 1986. Nucleotide sequence of pre-pro-von Willebrand factor cDNA. *Nucleic Acids Res.* 14:7125.

18. Shi, E., M. Kan, J. Xu, and W.L. McKeehan. 1991. 16-kilodalton heparin binding (fibroblast) growth factor type one appers in a stable 40-kilodalton complex after receptor-dependent internalization. *J. Biol. Chem.* 266:5774.
19. Catoretti, G., M.H.G. Becker, G. Key, M. Duchrow, C. Schlüter, J. Galle,  
5 and J. Gerdest. 1992. Monoclonal antibodies against recombinant parts of  
the Ki-67 antigen (MIB 1 and MIB 3) detect proliferating cells in microwave-  
processed formalin-fixed paraffin section. *J. of Pathol.* 168:357.

CLAIMS

1. An antibody directed against an FLT4 receptor tyrosine kinase.
2. A polyclonal antibody directed against an FLT4 receptor tyrosine  
5 kinase.
3. A monoclonal antibody directed against an FLT4 receptor tyrosine kinase.
- 10 4. The monoclonal antibody according to claim 3, wherein said antibody is anti-FLT4 monoclonal antibody 9D9F9.
5. A hybridoma cell line producing the monoclonal antibody according to claim 4.
- 15 6. The hybridoma cell line according to claim 5, wherein said hybridoma cell line is deposited as DSM ACC 2210.
7. A monoclonal antibody produced by the hybridoma cell line according  
20 to claim 6.
8. A detectably-labelled antibody according to claim 1.
9. A nucleic acid probe specific for DNA encoding human FLT4  
25 comprising the nucleic acid sequence of SEQ ID NO. 1 or fragments thereof which specifically bind/hybridize to DNA encoding human FLT4.
10. A method for detecting FLT4 in a biological sample;  
comprising the steps of  
30 a) exposing a sample suspected of containing FLT4 to a detectably-labelled probe according to claim 6;  
b) washing the sample; and  
c) detecting the presence of said detectably-labelled probe.

11. A method for detecting FLT4 in a biological sample,  
comprising the steps of

- a) exposing a sample suspected of containing FLT4 to a detectably-labelled anti-FLT4 antibody;
- 5       b) washing the sample; and
- c) detecting the presence of said detectably-labelled anti-FLT4 antibody in said sample.

12. A method for imaging lymphatic vessels in a tissue sample,  
10       comprising the steps of:

- a) applying a detectably labelled anti-FLT4 antibody to a site suspected of containing lymphatic vessels; and
- b) detecting said detectably-labelled anti-FLT4 antibody bound to said tissue.

15

13. A method for diagnosing diseases characterized by changes in lymphatic vessels and HEVs, comprising the steps of:

- a) obtaining a tissue sample from a patient suspected of having a disease characterized by changes in lymphatic cells and HEVs;
- 20       b) exposing said tissue sample to a detectably-labelled anti-FLT4 antibody;
- c) washing the said tissue sample; and
- d) detecting the presence of said detectably-labelled anti-FLT4 in said tissue sample.

25

14. A pharmaceutical composition comprising a therapeutically effective amount of an FLT4-binding compound in a pharmaceutically acceptable diluent, adjuvant or carrier, said composition being in a form suitable for injection.

30

15. The pharmaceutical composition according to claim 14, wherein the FLT4-binding compound comprises a detectably labelled anti-FLT4 antibody in a pharmaceutically acceptable diluent, adjuvant or carrier, said composition being in a form suitable for injection.

35



16. A method for detecting lymphatic vessels and HEVs in an organism, comprising the steps of

a) injecting a pharmaceutical composition according to claim 14 or 15, into said organism;

5        b) detecting the amount of said detectably labelled FLT4-binding compound bound to sites comprising lymphatic vessels or HEVs in said organism.

17. The method according to claim 16, wherein the lymphatic tissue to be  
10 detected is lymph node tissue.

18. A method for stimulating or antagonizing the function of FLT4 in an organism comprising the step of providing a pharmaceutical composition according to claim 14 or 15 in an amount sufficient to stimulate or block the  
15 FLT4-receptor.

19. The method according to claim 18, wherein the FLT4 receptor activity is associated with a disease selected from the group consisting of metastatic cancers, lymphomas, lymphangiomas, inflammation (chronic or acute),  
20 infections and immunological diseases.

1 / 8




Fig. 1A






Fig. 1B

SUBSTITUTE SHEET

2 / 8

*Fig. 2*

SUBSTITUTE SHEET



*Fig. 3*  
SUBSTITUTE SHEET






Fig. 4


*Fig. 5*



*Fig. 6*


SUBSTITUTE SHEET

7/8

*Fig. 7*

SUBSTITUTE SHEET





*Fig. 8*

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/FI 95/00337

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K16/28 C12N5/20 C12Q1/68 G01N33/577 A61K39/395  
A61K49/00 //C07K14/71

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N C12Q G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ONCOGENE, vol. 8, no. 11, November 1993 BASINGSTOKE, GB, pages 2931-2937, K. PAJUSOLA ET AL. 'Two human FLT4 receptor tyrosine kinase isoforms with distinct carboxy terminal tails are produced by alternative processing of primary transcripts.' see abstract see figure 1 see "material and methods"</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1,2,8-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 October 1995

Date of mailing of the international search report

16.11.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2220 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Nooij, F

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/FI 95/00337

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ONCOGENE, vol. 8, no. 5, May 1993 BASINGSTOKE, GB, pages 1233-1240, F. GALLAND ET AL. 'The FLT4 gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor.' cited in the application see abstract see figures 2,4,7 see "material and methods" ---	1,2,8-11
X	CANCER RESEARCH, vol. 52, no. 20, 15 October 1992 BALTIMORE, MD, USA, pages 5738-5743, K. PAJUSOLA ET AL. 'FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines.' cited in the application see abstract see figures 1,2 ---	9,10
X	THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 178, no. 6, 1 December 1993 NEW YORK, NY, USA, pages 2077-2088, A. KAIPAINEN ET AL. 'The related FLT4, FLT1, and KDR receptor tyrosine kinases show distinct expression patterns in human fetal endothelial cells.' cited in the application see abstract see page 2078, left column, line 54 - right column, line 1 ---	9,10
A	WO,A,94 10202 (GENENTECH, INC.) 11 May 1994 see examples see claims ---	1-8, 11-19
A	WO,A,93 14124 (HELSINKI UNIVERSITY HOLDING, LTD.) 22 July 1993 see claims see examples ---	1,2,8-19
	---	
	--- --	

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/FI 95/00337

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 92, no. 8, 11 April 1995 WASHINGTON, DC, USA, pages 3566-3570, A. KAIPAINEN ET AL. 'Expression of the fms-like tyrosine kinase 4 gene becomes restricted to lymphatic endothelium during development.' see abstract see "material and methods" -----</p>	9,10

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 95/00337

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:

Please see annex!

2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/FI 95/00337

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

REMARK: Although claims 16 and 17 are directed to a diagnostic method, and claims 18 and 19 to a medical treatment, practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/FI 95/00337

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9410202	11-05-94	AU-B- 2928992	24-05-94
		EP-A- 0666868	16-08-95
		FI-A- 951987	26-04-95
		NO-A- 951609	27-04-95
-----			
WO-A-9314124	22-07-93	AU-B- 3353293	03-08-93
		CA-A- 2127540	22-07-93
		EP-A- 0620826	26-10-94
		FI-A- 943275	11-07-94
		JP-T- 7506242	13-07-95
-----			